University of California

Cooperative Extension-Sacramento County

Agriculture and Natural Resources

4145 Branch Center Road, Sacramento, CA 95827-3823 (916) 875-6913 Office • (916) 875-6233 Fax Website: sacmg.ucanr.edu

Garden Notes

GN 154

SOIL TEMPERATURE CONDITIONS FOR VEGETABLE SEED GERMINATION

Many internal and environmental conditions influence seed germination: condition of the seed, presence of water, sufficient air, soil conditions, and temperature. The chart below lists the optimum soil temperatures at which they do best. This does not mean your seeds will not germinate at higher or lower temperatures, but greater success will be achieved if optimum conditions are provided. Using a probe-type thermometer is an accurate method for testing soil temperatures prior to planting seeds and small plants.

Crops	Minimum (°F)	Optimum range (°F)	Maximum (°F)	
Asparagus	50	75-85	95	
Beans, Lima	60	75-85	85	
Beans, Snap	60	75-85	95	
Beets	40	65-85	95	
Broccoli	40	60-85	95	
Cabbage	40	60-85	95	
Carrots	40 65-85		95	
Cauliflower	40 65-85		95	
Celery	40	*	*	
Chard, Swiss	40	40 65-85		
Corn	50	65-95	105	
Cucumbers	60	65-95	105	
Eggplant	60	75-85	95	
Garlic	32	65-85	95	
Leeks	32	65-85	95	
Lettuce	32	60-75	85	
Muskmelons (Cantaloupe)	60	75-85	105	
Okra	60	85-95	105	
Onions	32	65-85	95	
Parsley	40	65-85	95	
Parsnips	32	65-75	85	
Peas	40	65-75	85	
Peppers	60	65-75	95	
Pumpkins	60	85-95	105	
Radishes	40	65-85	95	
Spinach	32	65-75	75	
Squash	60	85-95	105	
Tomatoes	50	65-85	95	
Turnips	40	60-95	105	
Watermelons	60	75-95	105	

Source: *California Master Gardener Handbook*, 2nd edition, Regents of the University of California, Division of Agriculture and Natural Resources, Publication 3382 (Table 5.2, page 114).

*Note: Celery requires diffuse light and a night temperature from 10° to 15°F lower than the day temperature for good germination. Optimal conditions are 85°F day, 70°F night with diffuse light and high moisture.

It is the policy of the University of California (UC) and the UC Division of Agriculture & Natural Resources not to engage in discrimination against or harassment of any person in any of its programs or activities. (Complete nondiscrimination policy statement can be found at http://ucanr.edu/sites/anrstaff/files/215244.pdf). Inquiries regarding ANR's nondiscrimination policies may be directed to John Sims, Affirmative Action Compliance Officer/Title IX Officer, University of California, Agriculture and Natural Resources, 2801 Second Street, Davis, CA 95618, (530) 750-1397.

Agriculture • Nutrition, Family and Consumer Sciences • Master Food Preservers • 4-H Youth Development • Horticulture • Master Gardeners U.S. Department of Agriculture, University of California, and the Counties of Sacramento, Solano, and Yolo cooperating

NUMBER OF DAYS FOR VEGETABLE SEEDS TO EMERGE AT DIFFERENT TEMPERATURES

This chart shows the range of temperatures at which seeds will germinate. Note that temperatures higher or lower than the optimum soil temperature (as noted on the previous chart) will result in more days before germination occurs, or if temperatures are too extreme, germination may not result at all. For example, the optimum temperature range for tomatoes is 65° to 85°F (see previous chart). Within that range, it takes approximately 6 to 8 days before seeds germinate. Tomato seeds may still germinate at 50°F, but it will take over 40 days, and there will probably be no germination if the soil temperature is 104°F.

Crops	32°F	41°F	50°F	59°F	68°F	77°F	86°F	95°F	104°F
Asparagus	0.0	0.0	52.8	24.0	14.6	10.3	11.5	19.3	28.4
Beans, Lima			0.0	30.5	17.6	6.5	6.7	0.0	
Beans, Snap	0.0	0.0	0.0	16.1	11.4	8.1	6.4	6.2	0.0
Beets		42.0	16.7	9.7	6.2	5.0	4.5	4.6	
Cabbage	-		14.6	8.7	5.8	4.5	3.5		
Carrots	0.0	50.6	17.3	10.1	6.9	6.2	6.0	8.6	0.0
Cauliflower			19.5	9.9	6.2	5.2	4.7		
Celery	0.0	41.0	16.0	12.0	7.0	0.0	0.0	0.0	
Corn, Sweet	0.0	0.0	21.6	12.4	6.9	4.0	3.7	3.4	0.0
Cucumbers	0.0	0.0	0.0	13.0	6.2	4.0	3.1	3.0	
Eggplant	0.0				13.1	8.1	5.3		
Lettuce	49.0	14.9	7.0	3.9	2.6	2.2	2.6	0.0	0.0
Muskmelons	-				8.4	4.0	3.1		
Okra	0.0	0.0	0.0	27.2	17.4	12.5	6.8	6.4	6.7
Onions	135.8	30.6	13.4	7.1	4.6	3.6	3.9	12.5	0.0
Parsley			29.0	17.0	14.0	13.0	12.3		
Parsnips	171.7	56.7	26.6	19.3	13.6	14.9	31.6	0.0	0.0
Peas		36.0	13.5	9.4	7.5	6.2	5.9		
Peppers	0.0	0.0	0.0	25.0	12.5	8.4	7.6	8.8	0.0
Radishes	0.0	29.0	11.2	6.3	4.2	3.5	3.0		
Spinach	62.6	22.5	11.7	6.9	5.7	5.1	6.4	0.0	0.0
Tomatoes	0.0	0.0	42.9	13.6	8.2	5.9	5.9	9.2	0.0
Turnips	0.0	0.0	5.2	3.0	1.9	1.4	1.1	1.2	2.5
Watermelons		0.0			11.8	4.7	3.5	3.0	

Sources: J.F. Harrington, Department of Vegetable Crops, University of California, Davis, Agricultural Extension Leaflet, 1954

"The New Seed-Starting Handbook", Nancy Bubel, 1988, Rodale Press

Notes: 0.00 = little or no germination

--- = not tested

August 2017, updated. December 2015, revised and February 2013, written by UCCE Sacramento County Master Gardener Gail Pothour, reviewed by UCCE Master Gardeners Kim Brady, Dan Vierria, and Dave Vaughan. Edited by Judy McClure, UCCE Sacramento County Master Gardener Coordinator.